

Characterization of Spectral Entanglement of Spontaneous Parametric-Down Conversion Biphotons

G. Brida¹, V. Caricato^{1,2}, M. Genovese¹, M. Gramegna¹
¹*I.N.R.I.M.- Istituto Nazionale di Ricerca Metrologica, Turin, Italy*
²*Department of Physics, Politecnico of Turin, Turin, Italy*

M. V. Fedorov
A.M.Prokhorov General Physics Institute, Russian Academy of Science, Moscow, Russia

S. P. Kulik
Faculty of Physics, M.V.Lomonosov Moscow State University, Moscow, Russia
 (Dated: April 20, 2009)

We verified operational approach based on direct measurement of entanglement degree for bipartite systems. In particular spectral distributions of single counts and coincidence for pure biphoton states generated by train of short pump pulses have been measured and entanglement quantifier calculated. The approach gives upper bound of entanglement stored in total biphoton states, which can reach extremely high value up to $10^4 - 10^5$.

PACS numbers: 42.65.Lm, 03.65.Ud, 03.67.Mn

Recently entanglement became from an argument of debate on foundations on quantum mechanics a resource for developing quantum technologies [1] (as quantum communication, q-calculus, q-imaging, q-metrology, etc.). In this sense a precise and easy-implementable method for characterizing entanglement properties is a fundamental tool. There are several measures that quantify entanglement of quantum bipartite states both in discrete and continuous variables [2]. The most popular ones are Schmidt rank, entropy, concurrence, etc. From a theoretical point of view all these quantifiers can be easily evaluated from the density matrix describing the given state, thus there is no problem to find the meaning of a chosen entanglement measure for a certain density matrix. From an experimental side and following the same paradigm it would be necessary to perform quantum tomography procedure (either complete or reduced) over the state [3] under study and then to extract the entanglement measure from reconstructed density matrix. However this scheme is not optimal in the case of high-dimensional systems when number of measurements for complete reconstruction of the density matrix grows quadratically with dimension of Hilbert space. An alternative is exploiting particular links between entanglement and measurable characteristics of the state, just measuring some auxiliary parameters of the state. In particular the parameter R_q (Fedorov's ratio [4]), defined as the ratio of the single-particle and coincidence distributions widths in q -space, can be rather easily measured in contrast to all other entanglement quantifiers. R can be understood qualitatively from entropy approach to entanglement: high entanglement leads to the better knowledge about composite bipartite system (narrower coincidence distribution) and the worse knowledge about individual subsystem(s) (wider single-particle distribution).

Both clear physical meaning and operationability return R to an extremely useful tool for entanglement control. It has been proved [4] that for double-Gaussian bipartite states the parameter R coincides exactly with the Schmidt number K . Moreover their values remain quite close even for special classes of non-double-Gaussian wave functions like those describing Spontaneous Parametric Down-Conversion (SPDC). This approach can be used both for continuous and discrete variables, so the parameter R has rather universal sense: earlier it has been successfully applied for exposure strong entanglement anisotropy in spatial distributions of biphotons [5].

The present paper is devoted to verification of the operational entanglement quantifier R for two-photon states entangled in frequency domain as an efficient alternative to other quantifiers (for example in specific experiments visibility of interference pattern can provide some knowledge about entanglement [6]). Indeed such states belong to multi-dimensional Hilbert space and can possess extremely high entanglement degree (up to several hundreds) that makes its very perspective objects for quantum information and quantum communication.

Let us remind first the main features of the biphoton spectral wave function for SPDC with the type-I degenerate collinear phase matching and with a pulsed pump. The main general expression has the form [7, 8]

$$\Psi(\nu_1, \nu_2) \propto \exp\left(-\frac{(\nu_1 + \nu_2)^2 \tau^2}{8 \ln 2}\right) \times \text{sinc}\left\{\frac{L}{2c}\left[A(\nu_1 + \nu_2) - B\frac{(\nu_1 - \nu_2)^2}{\omega_p}\right]\right\}, \quad (1)$$

where τ is the pump-pulse duration, L is the length of the crystal, ν_1 and ν_2 are deviations of frequencies of the signal and idler photons $\omega_{1,2}$ from the central frequencies

all experiments were performed as a consecution of corresponding measurements of spectral distributions both in single counts and coincidences. To measure the coincidence distribution one of the two monochromators (by convention selecting idler wavelength) is fixed at the central wavelength of SPDC (795 nm) whilst the other (signal) scans in a range around this value. In order to check R dependence on the control parameter η the measurements have been repeated for crystals with different length (but with the same orientation), namely $L=10$ mm and $L=5$ mm.

First of all we have performed measurements with a 10 mm LiIO_3 crystal. To measure the coincidence distribution we have used monochromators with 0.2 nm resolution. A narrow peak with $\Delta\lambda_c = (0.29 \pm 0.03)$ nm has been obtained (Fig.3) while scanning the wavelength in signal channel. It is clearly seen that the pump width is 6.2 times larger than the coincidence spectrum.




FIG. 3: Normalized coincidence distribution for 10 mm length LiIO_3 sample. Points with error bars correspond to the measurements while the solid line shows the fitting by a Gaussian function with $\text{FWHM} = (0.29 \pm 0.03)$ nm.

The spectral distribution of single counts was performed using a monochromator in the transmission arm with a spectral resolution of 1 nm, finding a single counts spectral width $\Delta\lambda_s = (101 \pm 1)$ nm (Fig.4). The asymmetry of the right wing in the measured spectrum might be caused by falling spectral sensitivity of monochromator in the long-wavelength range.

Thus the distribution of single counts is 56 times larger than that of the pump, corresponding to an experimental ratio between widths of the two distributions $R_\omega = (349 \pm 43)$. The complete results are collected in the table I.

Then the same measurement has been performed with a crystal of 5 mm length obtaining $\Delta\lambda_s = (115 \pm 1)$ nm and $\Delta\lambda_c = (0.64 \pm 0.06)$ nm, corresponding to $R_\omega = (179 \pm 18)$. Considering that the pulse duration of the pump did not change we can observe that in this case the pump is 2.8 times greater than the coincidence spectrum whereas the single counts distribution is 64 times larger than the width of the pump. Corresponding results are




FIG. 4: Normalized single counts distribution for 10 mm length LiIO_3 sample.

TABLE I: Results for 10 mm LiIO_3 sample. The pump parameters τ and $\Delta\lambda_p$ are measured independently by using a cross-correlator (with first harmonic) and a spectrometer.

Theory Model	Experimental results
Pump	
$\lambda_p = 397.5 \text{ nm}$	$\lambda_p = 397.5 \pm 0.2 \text{ nm}$
$\Delta\lambda_p = 1.25 \text{ nm}$	$\Delta\lambda_p = (1.8 \pm 0.1) \text{ nm}$
$\tau = 186 \text{ fs}$	$\tau = 186 \pm 30 \text{ fs}$
Coincidence Distribution	
$\lambda_c = 795 \text{ nm}$	$\lambda_c = 795.0 \pm 0.2 \text{ nm}$
$\Delta\lambda_c = 0.32 \text{ nm}$	$\Delta\lambda_c = (0.29 \pm 0.03)$
Single Counts Distribution	
$\Delta\lambda_s = 100 \text{ nm}$	$\Delta\lambda_s = (101 \pm 1) \text{ nm}$
R -Quantifier	
$R_\omega \approx R_\lambda \approx 316$	$R_\omega \approx R_\lambda \approx (349 \pm 43)$

listed in the table II.

TABLE II: Results for 5 mm LiIO_3 sample

Theory Model	Experimental results
Coincidence Distribution	
$\Delta\lambda_c = 0.63 \text{ nm}$	$\Delta\lambda_c = (0.64 \pm 0.06)$
Single Counts Distribution	
$\Delta\lambda_s = 100 \text{ nm}$	$\Delta\lambda_s = (115 \pm 1) \text{ nm}$
R -Quantifier	
$R_\omega \approx R_\lambda \approx 158$	$R_\omega \approx R_\lambda \approx (179 \pm 18)$

Certainly the most comprehensive contribution to the experimental verification of the R -quantifier validity in frequency domain would be testing dependence R on the control parameter η through all available parameters. Agreement between the measured and predicted shapes of the curve shown at Fig.1 would confirm completely the adequacy of the R -quantifier approach. Looking at (3, 4) it is seen that the "working" parameters might be τ and L . However it is difficult to change the pump pulse dura-

tion in a wide range with the same laser (from hundreds of fs to dozens of ps) and pass through the minimum at the curve $R(\eta)$ varying this parameter only. That is why we took into account the fact of linear dependence of R on the sample length L through the coincidence distribution width (4). So doubling the sample length leads to doubling the entanglement degree for the short pump pulse regime of SPDC. This fact is clearly illustrated by the obtained results, demonstrating the validity of the conceptual scheme even if, unfortunately, subsequent decrease of the sample length reduces the biphoton flux too much for measuring widths $\Delta\lambda_s$ and reaching the minimum of $R(\eta)$.

One should mention that in some works (see, e.g., [10]) the 2D- photon distribution was measured and plotted in the (ω_1, ω_2) plane. Surprisingly that such 2D-distributions are very useful for making entanglement analysis of biphoton states in context of conditional pure state preparation, dispersion broadening [12, 13], etc. This would allow evaluating R -quantifier directly. We would like also to mention that a measurement of spectral width could also be obtained by transforming spectrum in time difference through a fiber [11].

Finally the question arises "how much entanglement can be stored in a given state". We remind that the method of measurement of R in frequency domain implies that other degrees of freedom of biphotons should be fixed. First off all we mean spatial distribution of photon pairs emitted by crystal, which must be illuminated by selecting particular narrow-angular segments with small pinholes. Vice versa, measurement of R -quantifier in spatial variables (R_{angle}) implies fixing frequencies of both photons, which is usually performed with narrow-band filters [5]. Therefore the total entanglement potentially stored in the SPDC state can be conventionally accepted as a product $R_{angle} * R_\omega$. On the one hand this is a good convention since quantifier R has the physical meaning of effective Hilbert space, so the total dimension of biphoton states in joint polarization-frequency-spatial space would be $R_{tot} \approx 2 * R_{angle} * R_\omega$. However we suggest to use this just as an upper bound of the stored entanglement taking into account the existing links between frequency and spatial degrees of freedom caused by phase matching. Speaking in terms of numbers this value can reach really huge amount $R_{tot} \approx 10^4 - 10^5$ [14]!

In conclusion we have verified the approach of Ref. [4] addressed to estimate the entanglement degree in

frequency domain for pure biphoton states generated by short pump pulses. An approach that can find widespread application in quantum technologies exploiting entanglement of biphotons. The generalization to the mixed two-photon states will be discussed elsewhere.

This work has been supported in part by MIUR (PRIN 2007FYETBY), Regione Piemonte (E14), "San Paolo foundation", NATO (CBP.NR.NRCL 983251) and RFBR (08-02-12091).

-
- [1] M. Genovese, Phys. Rep. 413, 3197 (2005).
 - [2] I. Bengtsson and K. Życzkowski, Geometry of Quantum States, Cambridge Univ. press (2006).
 - [3] A.I. Lvovsky et al, Phys. Rev. Lett. 87 (2001) 050402; A. Zavatta et al., Phys. Rev. A 70, 053821 (2004); A. Allevi et al., arXiv:0903.0104.
 - [4] M.V. Fedorov, M.A. Efremov, P.A. Volkov, and J.H. Eberly, J. Phys. B: At. Mol. Opt. Phys., **9**, S467 (2006).
 - [5] M.V. Fedorov, M.A. Efremov, P.A. Volkov, E.V. Moreva, S.S. Straupe, S.P. Kulik, Phys. Rev. Lett, **99**, 063901 (2007); Phys. Rev. A, **77**, 032336 (2008).
 - [6] Y. Kim et al., Phys. Rev. A 62 (2000) 043820; Y. Jeronimo-Moreno and A. B. U'Ren, Phys. Rev. A **79**, 033839 (2009).
 - [7] Yu.M. Mikhailova, P.A. Volkov, and M.V.Fedorov, Phys. Rev. A **78**, 062327 (2008).
 - [8] T.E. Keller and M.H. Rubin, Phys. Rev. A **56**, 1534 (1997).
 - [9] W. Mauerner and C. Silberhorn, (to be published in QCMC'08 Proceedings AIP, (2008).
 - [10] P.J. Mosley, J.S. Lundeen, B.J. Smith, P.Wasylczyk, A.B. U'Ren, Ch. Silberhorn, and I.A. Walmsley, Phys. Rev. Lett., **100**, 133601 (2008); Alejandra Valencia, Alessandro Cer, Xiaojuan Shi, Gabriel Molina-Terriza, and Juan P. Torres, Phys. Rev. Lett. 99, 243601 (2007); M. Avenhaus et al., arXiv 0810.0998.
 - [11] G. Brida, M. Chekhova, M. Genovese, M.Gramegna, L. Krivitsky. Phys. Rev. Lett. 96, 143601 (2006); G. Brida, M. Genovese, L. A.Krivitsky, M. V. Chekhova, Phys. Rev. A 75, 015801 (2007); M. Avenhaus, A. Eckstein, P. J. Mosley, C. Silberhorn, arXiv:0902.3364 (2008).
 - [12] S. Y. Baek, O. Kwon, and Y. H.Kim. Phys. Rev. A **77**, 013829 (2008); S. Y. Baek, O. Kwon, and Y. H.Kim. Phys. Rev. A **78**, 013816 (2008).
 - [13] R. Erdmann, D. Branning, W. Grice, and I. A. Walmsley. Phys. Rev. A **62**, 053810 (2000).
 - [14] Previous research [5] has demonstrated that extremely high entanglement degree can be accumulated in spatial variables of biphoton states ($R_{angle} \approx 300$).